Chronic Effects of Low pH on Some Physiological Aspects of Smoltification in Atlantic Salmon *(Salmo salar)*

C. E. Johnston, R. L. Saunders, E. B. Henderson,
P. R. Harmon and K. Davidson

Biological Station,
St. Andrews. N. B., E0G 2X0

August 1984

Canadian Technical Report of Fisheries and Aquatic Sciences No. 1294

84- 4961 /

Canadian Technical Report of Fisheries and Aquatic Sciences

These reports contain scientific and technical information that represents an important contribution to existing knowledge but which for some reason may not be appropriate for primary scientific (i.e. *Journal*) publication. Technical Reports are directed primarily towards a worldwide audience and have an international distribution. No restriction is placed on subject matter and the series reflects the broad interests and policies of the Department of Fisheries and Oceans, namely, fisheries management, technology and development, ocean sciences, and aquatic environments relevant to Canada.

Technical Reports may be cited as full publications. The correct citation appears above the abstract of each report. Each report will be abstracted in *Aquatic Sciences and Fisheries Abstracts* and will be indexed annually in the Department's index to scientific and technical publications.

Numbers 1-456 in this series were issued as Technical Reports of the Fisheries Research Board of Canada. Numbers 457-714 were issued as Department of the Environment, Fisheries and Marine Service, Research and Development Directorate Technical Reports. Numbers 715-924 were issued as Department of Fisheries and the Environment, Fisheries and Marine Service Technical Reports. The current series name was changed with report number 925.

Details on the availability of Technical Reports in hard copy may be obtained from the issuing establishment indicated on the front cover.

Rapport technique canadien des sciences halieutiques et aquatiques

Ces rapports contiennent des renseignements scientifiques et techniques qui constituent une contribution importante aux connaissances actuelles mais qui, pour une raison ou pour une autre, ne semblent pas appropriés pour la publication dans un journal scientifique. Il n'y a aucune restriction quant au sujet, de fait, la série reflète la vaste gamme des intérêts et des politiques du Ministère des Pêches et des Océans, notamment gestion des pêches, techniques et développement, sciences océaniques et environnements aquatiques, au Canada.

Les Rapports techniques peuvent être considérés comme des publications complètes. Le titre exact paraîtra au haut du résumé de chaque rapport, qui sera publié dans la revue *Aquatic Sciences and Fisheries Abstracts* et qui figurera dans l'index annuel des publications scientifiques et techniques du Ministère.

Les numéros 1-456 de cette série ont été publiés à titre de Rapports techniques de l'Office des recherches sur les pêcheries du Canada. Les numéros 457-714, à titre de Rapports techniques de la Direction générale de la recherche et du développement, Service des pêches et de la mer, ministère de l'Environnement. Les numéros 715-924 ont été publiés à titre de Rapports techniques du Service des pêches et de la mer, Ministère des Pêches et de l'Environnement. Le nom de la série a été modifié à partir du numéro 925.

La page couverture porte le nom de l'établissement auteur où l'on peut se procurer les rapports sous couverture cartonnée.

Canadian Technical Report of
Fisheries and Aquatic Sciences 1294

August 1984

CHRONIC EFFECTS OF LOW pH ON SOME PHYSIOLOGICAL ASPECTS OF SMOLTIFICATION IN ATLANTIC SALMON (*SALMO SALAR*)

by

C. E. Johnston[1], R. L. Saunders, E. B. Henderson, P. R. Harmon and K. Davidson[2]

Fisheries and Environmental Sciences
Fisheries Research Branch
Department of Fisheries and Oceans
Biological Station
St. Andrews, New Brunswick E0G 2X0

[1]Present address: Department of Biology, University of Prince Edward Island
Charlottetown, P.E.I. C1A 4P3
[2]Present address: Department of Fisheries and Oceans
Moncton, New Brunswick E1C 9B6

This is the one hundred and seventy-fourth Technical Report from the Biological Station, St. Andrews, N.B.

©Minister of Supply and Services Canada 1984
Cat. No. FS 97-6/1294E ISSN 0706-6457

Correct citation for this publication:

Johnston, C. E., R. L. Saunders, E. B. Henderson, P. R. Harmon, and K. Davidson, 1984. Chronic effects of low pH on some physiological aspects of smoltification in Atlantic salmon (Salmo salar). Can. Tech. Rep. Fish. Aquat. Sci. 1294: iii + 7 p.

ABSTRACT

Johnston, C. E., R. L. Saunders, E. B. Henderson, P. R. Harmon, and K. Davidson. 1984. Chronic effects of low pH on some physiological aspects of smoltification in Atlantic salmon (Salmo salar). Can. Tech. Rep. Fish. Aquat. Sci. 1294: iii + 7 p.

Atlantic salmon (Salmo salar) were chronically exposed for several months to acidified water to determine if mean pH levels of 4.7 and 4.9 prevent smoltification from proceeding normally. Physiological changes associated with smoltification developed normally when pH was above 4.9. When pH fell below 4.7, there was impairment of ionic regulatory mechanisms leading to excessive loss of electrolytes in fresh water. Failure of ionic regulation appeared to follow an inhibition of (Mg^{++})-ATPase and (Na^{+}, K^{+})-ATPase enzyme systems. Prolonged exposure to acid water allowed salmon to develop compensatory branchial mechanisms, including increased ATPase activity, that led to improved ionic regulation.

RÉSUMÉ

Johnston, C. E., R. L. Saunders, E. B. Henderson, P. R. Harmon, and K. Davidson. 1984. Chronic effects of low pH on some physiological aspects of smoltification in Atlantic salmon (Salmo salar). Can. Tech. Rep. Fish. Aquat. Sci. 1294: iii + 7 p.

Des saumons atlantiques (Salmo salar) ont été soumis à une exposition chronique de plusieurs mois à de l'eau acidifiée dans le but de déterminer si le processus normal de la smoltification était entravé à des pH moyens de 4,7 et 4,9. Les changements physiologiques associés à la smoltification se produisent normalement quand le pH est au-dessus de 4,9. Quand ce dernier baisse au-dessous de 4,7, il y a altération des mécanismes de régulation ionique, qui conduit à une perte excessive d'électrolytes en eau douce. Le manque de régulation ionique semble être la conséquence d'une inhibition des systèmes enzymatiques (Mg^{++})-ATPase et (Na^{+}, K^{+})-ATPase. Une exposition prolongée à de l'eau acide permet au saumon de développer des mécanismes branchiaux compensatoires, y compris une activité accrue de l'ATPase, qui résultent en une régulation ionique améliorée.

INTRODUCTION

Exposure of salmonids to pH conditions below 4.5 increases gill membrane permeability, mucus secretion from gills and skin, hydrogen ion uptake, blood acidosis, hematocrit, hemoglobin concentration, hemopoietic activity, respiratory and metabolic rates and loss of sodium, chloride and calcium ions from the plasma and body tissue (Flonka and Neff 1969; Packer and Dunson 1970; Vaala and Mitchell 1970; Hargis 1975; Daye and Garside 1976; Dively et al. 1977; Neville 1979a, b; Packer and Sunkin 1979; McDonald et al. 1980; Booth et al. 1981; Milligan and Wood 1982; Saunders et al. 1983). Other physiological consequences of acidification in salmonids are reduced active transport processes in chloride cells of the gill, inhibition of (Na^+,K^+)-ATPase activity and a decreased oxygen carrying capacity of the blood (Packer 1979; McDonald et al. 1983; Saunders et al. 1983).

Tolerance to low pH varies with species, developmental stage and length of exposure. Grande et al. (1979) reported that, of four salmonid species, brook trout (Salvelinus fontinalis) are more acid tolerant than brown trout (Salmo trutta), Atlantic salmon or rainbow trout (Salmo gairdneri). Eggs, alevins and fry of Atlantic salmon are the most sensitive developmental stages and many physiological and pathological alterations occur in the presence of high hydrogen ion concentrations (Daye and Garside 1977, 1979, 1980; Peterson et al. 1980; Haya and Waiwood 1981). Most insight into the toxic effects of low pH on salmonids has been gained through experiments involving acute exposure to acid, usually lasting less than 7 d. Only a few experiments have employed chronic acid exposures of longer duration, and most have dealt with egg or alevin stages (Spry et al. 1981; Wood and McDonald 1982). Although Atlantic salmon are one of the most pH sensitive and economically important species in Atlantic Canada, few chronic laboratory or field experiments have been conducted on juveniles (fry, parr and smolts) or adults returning from the sea. Knowledge of the toxic levels and of the impact of prolonged exposure to hydrogen ions on seasonally determined endogenous rhythms and on physiological and biochemical mechanisms still remains poorly elucidated.

In a recent long-term experiment with Atlantic salmon we reared parr in low pH environments (4.2-4.7) at constant 10°C and rising temperature to the smolt stage (Saunders et al. 1983). Salmon exposed to low pH during the period when they should have been undergoing parr-smolt transformation failed to develop tolerance to high salinity, failed to elevate branchial (Na^+,K^+)-ATPase activity, failed to maintain normal plasma Na^+, Cl^- and Ca^{++} concentrations and failed to maintain normal tissue moisture content and growth. Based on these results, the process of smoltification did not appear to proceed normally in the pH range 4.2-4.7 even though the fish survived. The present laboratory study was undertaken to define the threshold pH above which smoltification proceeds normally and to examine further the physiological changes that accompany chronic exposure to acid water during parr-smolt transformation.

MATERIALS AND METHODS

In mid-February 1982, 1500 yearling Atlantic salmon parr (12-15 cm fork length) were randomly distributed among six 1-m^3 Swedish style rearing tanks (250/tank) at the St. Andrews Biological Station. Water in these tanks was gradually heated from an ambient temperature of ca. 2°C in mid-February to 10°C by March 4, from which time it was maintained at this temperature for the remainder of the experiment. Flow rates were ca. 12 L/min. Initial sampling was conducted on March 5, after which three pH regimes were established in the six experimental tanks. Two control tanks used normal laboratory water with a mean ambient pH of 6.78 (range 6.55-6.99) and Na, Cl, Ca, Mg, K and SO_4 levels of 100, 100, 38, 15, 10 and 20 μmol/L, respectively. Three groups were held under low pH (mean pH 4.66, range 4.33-5.00) and the remaining group was at intermediate pH (mean pH 4.94, range 4.46-5.40). Experimental conditions were maintained by using the methodology outlined by Saunders et al. (1983). Although pH conditions were relatively constant, some fluctuations occurred during the study period; these fluctuations (expressed as mean pH over 5-d periods) are shown in Fig. 1.

The methodology involved in feeding, photoperiod regimes, length-weight measurements, moisture determinations, measurement of ATPase activity, salinity tolerance tests, and ion measurements follows those of Saunders et al. (1983) with the following changes: fish were fed to satiety at a rate of 2% body weight/d; blood samples were collected from the caudal artery, using 1-cc heparinized syringes; a Perkin-Elmer 303 atomic absorption spectrophotometer was used for analysis of Na^+ and Ca^{++} levels.

Statistical comparisons between means were made by using a Duncan's Multiple Range Test or a Student's t-test.



Fig. 1. Three pH regimes maintained during the experiment which ran from March 5-May 25, 1982.

RESULTS AND DISCUSSION

SURVIVAL AND GROWTH

Salmon exposed to the low pH treatment exhibited higher mortality (56.1%) during the experiment than those exposed to the intermediate and control pH regimes where mortality was 2.4% and 0%, respectively. Mortality in the low pH group followed a modal pattern that was directly related to excursions in pH level (Fig. 2). Initiation of the modes occurred when pH levels (calculated over 5-d period) dropped to less than 4.5. If the pH level was increased above this value, there was recovery, but the ameliorative effect of this increase was not immediate, creating a lag effect.

In the intermediate pH regime, there were six mortalities during the period May 14-16. The timing of this mortality coincided with late stages of smoltification in the control fish and may represent a lowered resistance to high concentrations of hydrogen ions and a progressive deterioration in health and condition owing to long-term exposure to acid water.

Growth of salmon exposed to control and intermediate pH regimes was similar and greater than those exposed to low pH (Fig. 3). Slower growth and the apparent decrease in size of salmon exposed to the low pH was probably caused by size-related mortality and a poor feeding response.

MOISTURE CONTENT AND CONDITION FACTOR

Total moisture content for salmon from the low pH regime, although slightly higher in April and early May and lower in late May than values for the other regimes, was not significantly different from those of the other pH regimes (Table 1).

The condition factor for salmon at low pH was significantly lower ($P < 0.05$) throughout the study period than those for the other regimes. Extremely low values were reached in late May. Such low condition factors suggest that salmon exposed to low pH lose body weight and body energy reserves at a rate greater than salmon at higher pH levels.

SALINITY TOLERANCE

Survival of salmon in initial salinity tolerance tests (conducted at 35 o/oo salinity) in March was 50% after 96 h for control pH conditions (Fig. 4); in subsequent tests, control fish had 80-100% survival. Survival of salmon in intermediate and low pH conditions was 60-70% and 0-30%, respectively, between mid-April and late May. These data clearly indicate that salinity tolerance of juvenile salmon was greatly impaired at low pH and to a lesser extent at pH 4.9.

BRANCHIAL ATPASE ACTIVITY

Branchial (Mg^{++})-ATPase activity was significantly reduced ($P < 0.05$) in salmon exposed to pH 4.7 (Fig. 5). In mid-April and early May (Mg^{++})-ATPase levels of the low pH group were significantly lower ($P < 0.05$) than those for either of the other regimes. By late May, however, (Mg^{++})-ATPase activity of the low pH group increased and was not significantly different from the other regimes. The pattern of (Mg^{++})-ATPase activity followed that of the plasma ions (Fig. 6, 7, 8).

Branchial (Na^{+}, K^{+})-ATPase activity was significantly lower during April and May ($P < 0.01$) in salmon exposed to the low pH regime than to the intermediate or control regimes (Fig. 5). Salmon exposed to the intermediate pH regime also had significantly lower ($P < 0.05$) ATPase levels than the controls during May. These results are in agreement with those of the salinity tolerance tests, indicating marked physiological impairment of osmoregulatory function in acid-stressed juvenile salmon in salt water. The marked decrease in ATPase activity and salinity tolerance in the controls between May 6th and 21st suggests that the smoltification process may have peaked in early May and that desmoltification commenced thereafter.

PLASMA ION CONCENTRATIONS

Mean plasma Na^{+}, Cl^{-} and Ca^{++} ion concentrations varied seasonally and in magnitude according

Table 1. Mean moisture content (expressed as a percentage of total wet body weight) and condition factor (CF) for Atlantic salmon parr-smolts exposed to three different pH regimes.

Sampling date	pH regime: Low (%)	Low (CF)	Intermediate (%)	Intermediate (CF)	Control (%)	Control (CF)
1982						
March 3	-		-		72.4	1.02
April 19	75.5	0.91	74.7	0.97	74.6	0.94
May 10	75.6	0.88	73.9	0.92	74.8	0.92
May 25	73.6	0.81	73.1	0.90	75.1	0.92



Fig. 2. Mortalities that occurred in Atlantic salmon populations maintained at low (4.66) pH levels during the period March 5-May 26, 1982.



Fig. 3. Fork lengths of Atlantic salmon reared at control (6.78), intermediate (4.94), and low (4.66) pH levels.



Fig. 4. Salinity tolerance of Atlantic salmon reared at control (6.78), intermediate (4.94), and low (4.66) pH levels. Salinity tolerance expressed as percent survival after 96 h in 35 o/oo seawater.



Fig. 5. Mean (Mg^{++})- and (Na^{+},K^{+})-ATPase activity levels (± SE in Atlantic salmon reared at control (6.78), intermediate (4.94), and low (4.66) pH levels. ATPase activity expressed as umole $P_i \cdot mg\ protein^{-1} \cdot h^{-1}$.



Fig. 6. Mean plasma Na^{+} levels (± SE) in Atlantic salmon reared at control (6.78), intermediate (4.94), and low (4.66) pH levels.



Fig. 8. Mean plasma Ca^{++} levels (± SE) in Atlantic salmon reared at control (6.78), intermediate (4.94), and low (4.66) pH levels.



Fig. 7. Mean plasma Cl^{-} levels (± SE) in Atlantic salmon reared at control (6.78), intermediate (4.94), and low (4.66) pH levels.

to the pH regime (Fig. 6, 7, 8). Except for the control salmon, the pattern of seasonal change in plasma ion concentrations for each of the other regimes was similar, consisting of an initial decline followed by a rise in ion concentrations. The greatest decline in ion concentrations occurred in salmon from the low pH regime. Levels of all ions in April and early May were significantly lower ($P < 0.05$) in salmon reared at low pH than in those at intermediate and control pH. By late May, those surviving in the low pH regime had plasma ion concentrations not significantly different from those of the other regimes. This rise in ion levels to near normal values in the low pH regime may reflect the slightly higher pH conditions at the time of sampling in early and late May (pH 4.53-4.63) compared with mid-April (pH 4.49-4.51), or it may reflect some form of branchial compensation in which plasma ion losses are reduced and active uptake of ions from the water is increased.

GENERAL DISCUSSION

The effect of long-term acid exposure on smoltification of Atlantic salmon has been described to some extent already by Saunders et al. (1983). However, in that experiment, it was not possible to demonstrate the level of pH above which smoltification would proceed normally. In this experiment, we have compared the effect of three pH regimes considered to be above and below the threshold level for smoltification.

Although pH below 4.7 did not appear to influence body silvering, the development of marginal fin pigmentation or the process of scale loss in smolts, it did have a marked effect on growth and condition factor. Growth data for the low pH regime are not suitable for statistical comparisons because of the high mortality. However, the data do suggest poorer growth in the low pH condition. Poor feeding response and a heightened level of swimming activity appear to be the main causes of this poor growth.

The rate of decline in condition factor for salmon exposed to low pH was much greater than rates for salmon in either of the other regimes. These changes in condition are probably associated with weight loss. The early depletion of lipid reserves and lower condition factor were probably caused by increased energy demands for muscle contraction, increased cardiovascular activity, and increases in other body functions affected by exposure to low pH.

These data also clearly demonstrate that physiological rhythms and mechanisms associated with ionic regulation are impaired when salmon are chronically exposed to pH conditions below 4.7. Smoltification appears to proceed when pH is at or above 4.9, since loss of plasma ions, reduced condition, impairment of ATPase activity and salinity tolerance are greatly attenuated.

The physiological mechanism of acid toxicity for chronically exposed juvenile Atlantic salmon during the time when they should be smolting appears to be associated with failure of ionic regulation in fresh water and sea water. Death in fresh water coincided with low plasma Na^+, Cl^- and Ca^{++} levels, and loss of body condition. Death in sea water of 35 o/oo salinity coincided with the high plasma Na^+ and Cl^- concentrations, increased hematocrit and low branchial (Na^+,K^+)-ATPase activity (Saunders levels and ATPase activity are much higher and close et al. 1983). When pH is 4.9 or higher, plasma ion to controls. Few salmon died in these conditions.

Loss of ions from the plasma of salmonids exposed to low pH conditions in these chronic experiments may be owing to several different actions. Ion losses could be caused by an increase in diffusive permeability of the branchial epithelium, inhibition of exchange diffusion mechanisms involving Na^+, H^+ and NH_4^+ or Cl^- and HCO_3^-, inhibition of active transport pumps associated with the branchial chloride cell, or to some combination of these actions (Wood and McDonald 1982). Present data suggest that ion loss may be caused by a failure of freshwater and saltwater ion transport mechanisms driven by (Mg^{++})-ATPase or (Na^+,K^+)-ATPase enzyme systems. In fresh water, a decline in (Mg^{++}) and (Na^+,K^+)-ATPase activity coincided with declines in all plasma ion concentrations. Conversely, high (Mg^{++})-ATPase activity in fresh water always coincided with elevated plasma ion concentrations. Elevated (Na^+,K^+)-ATPase activity coincided with better survival in seawater and better regulation of Na^+ and Cl^- (Saunders et. al. 1983).

It has been suggested that an increase in branchial ATPase activity is directly related to the excretion of sodium and chloride by the gill in salt water or to the reabsorption of these ions in fresh water (Epstein et al. 1967; Kamiya and Utida 1968, 1969; Pickford et al. 1970; Johnson 1973). Although the role played by (Mg^{++})-ATPase and (Na^+,K^+)-ATPase enzymes in ion transport across the epithelial tissues of the gill remains speculative, data indicate that the pattern of ATPase activity follows the pattern of plasma Na^+ and Cl^- fluctuations. McDonald et al. (1983), in an experiment with rainbow trout, suggested that losses of Na^+ and Cl^- across the gills were caused by a pronounced inhibition of active transport mechanisms. The effect of hydrogen ions on the branchial ion transport system, however, is not clearly understood. Judging from the marked decline in (Mg^{++})-ATPase and (Na^+-K^+)-ATPase activity in low pH conditions, our data suggest that failure of the energy producing component in the chloride cells or in the transport mechanism is the underlying reason for failure of the ionic regulatory system.

Failure of active transport mechanisms to replace or remove plasma ions may not be the only cause of abnormal plasma ion concentrations. Changes in the intercellular cementing agents restricting the paracellular diffusion channels, changes in the permeability of the apical membranes of chloride cells or altered exchange diffusion rates could also lead to reductions in plasma ion concentrations (McDonald 1983). It is not possible to state which of these is most affected by low pH conditions.

In late May, salmon exposed to low pH conditions showed elevated (Mg^{++})-ATPase activity and near normal plasma ion concentrations in fresh water. This change in physiological response may be associated in some way with the smoltification process or with the development of acid tolerance that is independent of smolting. Some degree of acid resistance may exist, or develop, in survivors of exposure to low pH. Several studies have shown improved resistance of eggs and alevins of brook

trout and of juvenile brown trout during long-term studies (Trojnar 1977; McWilliams 1980). There is evidence also that elevated hormonal levels associated with smoltification could play an important role in stimulating branchial ATPase activity. Folmar and Dickhoff (1979) and Gallis et al. (1979) reported that gill ATPase activity was influenced by plasma thyroxine and cortisol levels. Since both of these hormones show seasonal changes, it is possible that increased activity of these hormones is responsible for this change in acid tolerance. Further investigations are needed to describe the nature of the structural and functional interactions that develop in the chloride cells of Atlantic salmon chronically exposed to low pH during smoltification.

In conclusion, we suggest that the pH level at which smoltification will proceed normally is at or above 4.9. If pH falls below this level, some impairment of ionic-regulatory mechanisms occurs, leading to excessive loss or gain of electrolytes in fresh water and sea water, respectively. The primary causes of ionic-regulatory failure in this study appear to be associated with inhibition of (Mg^{++})-ATPase and (Na^{+},K^{+})-ATPase systems and possibly altered membrane permeability in fresh water and salt water. Following prolonged exposure to acid media, some smolting salmon appear to develop compensatory branchial mechanisms and improved ionic regulation. The nature of the compensatory mechanisms is not understood.

ACKNOWLEDGMENTS

We thank Mr. Tim Birt for assisting with the collection of tissue samples and ATPase measurements. Drs. R. H. Peterson and G. L. Lacroix reviewed the manuscript. Messrs. P. W. G. McMullon and F. Cunningham prepared the illustrations; Mrs. E. Blanchard and Mrs. B. Fawkes typed the manuscript.

Part of this study was supported by grants received by C. E. Johnston from the National Sciences and Engineering Research Council of Canada, and the University of Prince Edward Island Research Committee.

REFERENCES

Booth, J. H., G. F. Jansz, and G. F. Holeton. 1981. Cl^{-}, K^{-}, and acid-base balance in rainbow trout during exposure to, and recovery from, sublethal environmental acidification. Can. J. Zool. 60: 1123-1130.

Daye, P. G., and E. T. Garside. 1976. Histopathologic changes in surficial tissues of brook trout, Salvelinus fontinalis (Mitchill), exposed to acute and chronic levels of pH. Can. J. Zool. 54: 2140-2155.

1977. Lower lethal levels of pH for embryos and alevins of Atlantic salmon, Salmo salar L. Can. J. Zool. 55: 1504-1508.

1979. Development and survival of embryos and alevins of the Atlantic salmon, Salmo salar L., continuously exposed to acidic levels of pH, from fertilization. Can. J. Zool. 57: 1713-1718.

1980. Structural alternations in embryos and alevins of the Atlantic salmon, Salmo salar L., induced by continuous or short-term exposure to acidic levels of pH. Can. J. Zool. 58: 27-43.

Dively, J., J. Mudge, W. Neff, and A. Anthony. 1977. Blood Po_2, Pco_2 and pH changes in brook trout (Salvelinus fontinalis) exposed to sublethal levels of acidity. Comp. Biochem. Physiol. 57A: 347-351.

Epstein, F. H., A. I. Katz, and G. E. Pickford. 1967. Sodium and potassium activated adenosine triphosphatase of gills: Role in adaptation of teleosts to salt water. Science 156: 1245-1247.

Folmar, L. C., and W. W. Dickhoff. 1979. Plasma thyroxine and gill Na^{+}-K^{+}-ATPase changes during seawater acclimation of coho salmon, Oncorhynchus kisutch. Comp. Biochem. Physiol. 63A: 329-332.

Gallis, J. L., F. Belloc, P. Lassere, and J. Boisseau. 1979. Freshwater adaptation in the euryhaline teleost, Chelon labrosus. II. Effects of continuance of adaptation cortisol treatment, and environmental calcium on water flux in isolated gill. Gen. Comp. Endocrinol. 38: 11-20.

Grande, M., I. P. Muniz, and S. Andersen. 1979. The relative tolerance of some salmonids to acid waters. Verh. Internat. Verein. Limnol. 20: 2076-2084.

Hargis, J. R. 1975. Ventilation and metabolic rate of young rainbow trout (Salmo gairdneri) exposed to sublethal environment pH. J. Exp. Zool. 196: 39-43.

Haya, K., and B. A. Waiwood. 1981. Acid pH and chorionase activity of Atlantic salmon (Salmo salar) eggs. Bull. Environmn. Contam. Toxicol. 27: 7-12.

Johnson, D. W. 1973. Endocrine control of hydromineral balance in teleosts. Amer. Zool. 13: 799-818.

Kamiya, M., and S. Utida. 1968. Changes in activity of Na-K activated adenosine triphosphatase in gills during adaptations of the Japanese eel to sea water. Comp. Biochem. Physiol. 26: 675-685.

1969. Sodium-potassium activated adenosine triphosphatase activity in gills of freshwater, marine and euryhaline teleosts. Comp. Biochem. Physiol. 31: 671-674.

McDonald, D. G. 1983. The effects of H^{+} upon the gills of freshwater fish. Can. J. Zool. 61: 691-703.

McDonald, D. G., H. Hobe, and C. M. Wood. 1980. The influence of calcium on the physiological responses of the rainbow trout, Salmo gairdneri, to low environmental pH. J. Exp. Biol. 88: 109-131.

McDonald, D. G., R. L. Walker, and P. R. H. Wilkes. 1983. The interaction of environmental calcium and low pH on the physiology of the rainbow trout, Salmo gairdneri. II. Branchial ionoregulatory mechanisms. J. Exp. Biol. 102: 141-155.

McWilliams, P. G. 1980. Acclimation to an acid medium in the brown trout Salmo trutta. J. Exp. Biol. 88: 269-280.

Milligan, C. L., and C. M. Wood. 1982. Disturbances in haematology, fluid volume distribution and circulatory function associated with low environmental pH in the rainbow trout, Salmo gairdneri. J. Exp. Biol. 99: 397-415.

Neville, C. M. 1979a. Ventilatory response of rainbow trout (Salmo gairdneri) to increased H^+ concentration in blood and water. Comp. Biochem. Physiol. 63A: 373-376.

1979b. Sublethal effects of environmental acidification on rainbow trout (Salmo gairdneri). J. Fish. Res. Board Can. 36: 84-87.

Packer, R. K. 1979. Acid-base balance and gas exchange in brook trout (Salvelinus fontinalis) exposed to acidic environments. J. Exp. Biol. 79: 127-134.

Packer, R. K., and W. A. Dunson. 1970. Effects of low environmental pH on blood pH and sodium balance in brook trout. J. Exp. Zool. 174: 65-71.

Packer, R. K., and A. L. Sunkin. 1979. Blood acid-base balance in brook trout (Salvelinus fontinalis). J. Exp. Biol. 79: 115-126.

Peterson, R. H., P. G. Daye, and J. L. Metcalfe. 1980. Inhibition of Atlantic salmon hatching at low pH. Can. J. Fish. Aquat. Sci. 37: 770-774.

Pickford, G. E., R. W. Griffith, J. Torrete, E. Hendler, and F. H. Epstein. 1970. Branchial reduction and renal stimulation of $(Na^+ + K^+)$ ATPase by prolactin in hypophysectomized killifish in fresh water. Nature (London) 226: 378-379.

Plonka, A., and W. Neff. 1969. Mucopolysaccharide histochemistry of gill epithelial secretions in brook trout exposed to acid pH. Proceedings of the Pennsylvania Academy of Sciences 43: 53-56.

Saunders, R. L., E. B. Henderson, P. R. Harmon, C. E. Johnston, and J. G. Eales. 1983. Effects of low environmental pH on smolting of Atlantic salmon (Salmo salar). Can. J. Fish. Aquat. Sci. 40: 1203-1211.

Spry, D. J., C. M. Wood, and P. V. Hodson. 1981. The effects of environmental acid on freshwater fish, with particular reference to the softwater lakes in Ontario and the modifying effects of heavy metals in the environment. A literature review. Can. Tech. Rep. Fish. Aquat. Sci. 999: 144 p.

Trojnar, J. R. 1977. Egg hatchability and tolerance of brook trout (Salvelinus fontinalis) fry at low pH. J. Fish. Res. Board Can. 34: 574-579.

Vaala, S., and R. Mitchell. 1970. Blood oxygen-tension changes in acid-exposed brook trout. Proceedings of the Pennsylvania Academy of Sciences 44: 41-44.

Wood, C. M., and D. G. McDonald. 1982. Physiological mechanisms of acid toxicity to fish, p. 197-226. In The Proceedings of the Acid Rain Fisheries Symposium. Am. Fish. Soc., August 1981, Ithaca, New York.